

ERRATUM TO “FROM LOOP GROUPS TO 2-GROUPS”

JOHN C. BAEZ, ALISSA S. CRANS, URS SCHREIBER AND DANNY STEVENSON

Abstract

There were a number of sign errors in our paper “From loop groups to 2-groups” [*Homology Homotopy Appl.* **9** (2007), 101–135]. Here we explain how to correct those errors.

The following sign corrections to our paper [BCSS] make the article consistent with [BC] and [LM], in particular the definition of morphisms between 2-term L_∞ -algebras. The definition given in [BC], Definition 4.3.4, was independently checked to be consistent with the one suggested in [LM, Remark 5.3] by Kevin van Helden. Since the main theorem of the paper involves several such morphisms, a number of signs in the data of some L_∞ -algebra morphisms need to be changed. Moreover, the definition of the crossed module action α in Proposition 2.4 led to an inconsistency independent of the L_∞ -algebra material, found by David Michael Roberts. The corrected sign is self-consistent, as well as agreeing with the rest of the paper. There were additionally some typos in an earlier paper [MS] leading to some innocuous sign errors that do not impact the other calculations.

All the corrections here have been made in the current arXiv version of our paper [BCSS]. All of our calculations have been independently checked by Rist, Saemann and Wolf [RSW], as well as by Roberts.

- The big commutative diagram in Definition 2.2 should be replaced by

$$\begin{array}{ccc}
 [[F_0(x), F_0(y)], F_0(z)] & \xrightarrow{J_{F_0(x), F_0(y), F_0(z)}} & [F_0(x), [F_0(y), F_0(z)]] + [[F_0(x), F_0(z)], F_0(y)] \\
 \downarrow [F_2, 1] & & \downarrow [1, F_2] + [F_2, 1] \\
 [F_0[x, y], F_0(z)] & & [F_0(x), F_0[y, z]] + [F_0[x, z], F_0(y)] \\
 \downarrow F_2 & & \downarrow F_2 + F_2 \\
 F_0[[x, y], z] & \xrightarrow{F(J_{x, y, z})} & F_0[x, [y, z]] + F_0[[x, z], y]
 \end{array}$$

2020 Mathematics Subject Classification: 22E67

Article available at <http://dx.doi.org/10.4310/HHA.????v??n??a?>

Copyright © ????, John C. Baez, Alissa S. Crans, Urs Schreiber and Danny Stevenson. Permission to copy for private use granted.

- Equation (5) needs to be replaced by

$$\begin{aligned} \phi_1(l_3(x, y, z)) - l_3(\phi_0(x), \phi_0(y), \phi_0(z)) = \\ \phi_2(x, l_2(y, z)) + \phi_2(y, l_2(z, x)) + \phi_2(z, l_2(x, y)) + \\ l_2(\phi_0(x), \phi_2(y, z)) + l_2(\phi_0(y), \phi_2(z, x)) + l_2(\phi_0(z), \phi_2(x, y)) \end{aligned}$$

- The definition of κ in equation (12) should be

$$\kappa(f, g) = \exp\left(-2ik \int_0^{2\pi} \int_0^{2\pi} \langle f(t)^{-1} f'(t), g'(\theta)g(\theta)^{-1} \rangle d\theta dt\right),$$

correcting a typo in [MS].

- The definition of the normal subgroup N below equation (12) should read

Let N be the subset of $P_0\Omega G \times U(1)$ consisting of pairs (γ, z) such that $\gamma: [0, 2\pi] \rightarrow \Omega G$ is a loop based at $1 \in \Omega G$ and

$$z = \exp\left(ik \int_{D_\gamma} \omega\right)$$

where D_γ is any disk in ΩG with γ as its boundary.

for consistency with the definition of κ above.

- The definition of $d\alpha$ in the statement of Proposition 3.1 should be

$$d\alpha(p)(\ell, c) = ([p, \ell], 2k \int_0^{2\pi} \langle \ell(\theta), p'(\theta) \rangle d\theta).$$

- The definition of β_p in the proof of Proposition 3.1 should be

$$\beta_p(\xi) = 2 \int_0^{2\pi} \langle \xi(\theta), p(\theta)^{-1} p'(\theta) \rangle d\theta.$$

- The definition of ϕ_2 in the statement of Lemma 5.4 should be

$$\phi_2(p_1, p_2) = k \int_0^{2\pi} (\langle p_2, p_1' \rangle - \langle p_2', p_1 \rangle) d\theta$$

- The definition of λ_2 in the proof of Lemma 5.5 should be

$$\lambda_2(\ell_1, \ell_2) = (0, 2k \int_0^{2\pi} \langle \ell_1, \ell_2' \rangle d\theta)$$

With these changes, the calculations in the proofs all go through, leaving the results unchanged.

References

- [BC] J. Baez and A. S. Crans, Higher-dimensional algebra VI: Lie 2-algebras, *Theor. Appl. Categ.* **12** (2004), 492–528. Also available as [arXiv:math/0307263](https://arxiv.org/abs/math/0307263).
- [BCSS] J. C. Baez, A. S. Crans, U. Schreiber and D. Stevenson, From loop groups to 2-groups, *Homology Homotopy Appl.* **9** (2007), 101–135. Corrected version available as [arXiv:math/0504123v3](https://arxiv.org/abs/math/0504123v3).

- [**LM**] T. Lada and M. Markl, Strongly homotopy Lie algebras, *Commun. Alg.* **6** (1995), 2147–2161. Also available as [arXiv:hep-th/9406095](https://arxiv.org/abs/hep-th/9406095).
- [**MS**] M. K. Murray and D. Stevenson, Higgs fields, bundle gerbes and string structures, *Commun. Math. Phys.* **243** (2003), 541–555. Also available as [arXiv:math/0106179](https://arxiv.org/abs/math/0106179).
- [**RSW**] D. Rist, C. Saemann and M. Wolf, Explicit non-abelian gerbes with connections. Available as [arXiv:2203.00092](https://arxiv.org/abs/2203.00092).

John C. Baez baez@math.ucr.edu

Department of Mathematics, University of California at Riverside, U.S.A

Alissa S. Crans acrans@lmu.edu

Department of Mathematics, Loyola Marymount University, U.S.A.

Urs Schreiber urs.schreiber@gmail.com

Mathematics Division, New York University in Abu Dhabi, Abu Dhabi

Danny Stevenson daniel.stevenson@adelaide.edu.au

Mathematical Sciences, The University of Adelaide, Australia